

Weakly nonlinear dynamics and the σ_8 parameter

Roman Juszkiewicz^{a,b}, Hume A. Feldman^{c,d,e}, J. N. Fry^f, Andrew H. Jaffe^e

^a*Institute of Astronomy, 65-516 Zielona Góra*

^b*Copernicus Astronomical Center, 00-716 Warsaw, Poland*

^c*Department of Physics and Astronomy, University of Kansas, Lawrence KS 66045, USA*

^d*Department of Physics and Astronomy, University College London, London, WC1E 6BT, UK*

^e*Astrophysics, Blackett Laboratory, Imperial College, London SW7 2AZ, UK*

^f*Department of Physics, University of Florida, Gainesville FL 32611-8440, USA*

Email: roman@camk.edu.pl, feldman@ku.edu, fry@phys.ufl.edu, a.jaffe@imperial.ac.uk

ABSTRACT

The amplitude of cosmological density fluctuations, σ_8 , has been studied and estimated by analysing many cosmological observations. The values of the estimates vary considerably between the various probes. However, different estimators probe the value of σ_8 in different cosmological scales and do not take into account the nonlinear evolution of the parameter at late times. We show that estimates of the amplitude of cosmological density fluctuations derived from cosmic flows are systematically higher than those inferred at early epochs from the CMB because of nonlinear evolution at later times. We discuss the past and future evolution of linear and nonlinear perturbations, derive corrections to the value of σ_8 and compare amplitudes after accounting for these differences.

Subject headings: cosmic flows, cosmological parameters from LSS, cosmological parameters from CMBR, galaxy dynamics

1. Introduction

The σ_8 parameter is one of the most important and least well known parameters in cosmology. It is a convenient measure of the degree of the inhomogeneity of the Universe. It is the rms matter density contrast in a sphere with a comoving radius of $8h^{-1}$ Mpc at present, where h is the usual dimensionless Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$. The original motivation for the use of this parameter was the need to define the clustering amplitude: the variance in the number density of optical galaxies in $8h^{-1}$ Mpc spheres, σ_{gal}^2 , was observed to be unity (Davis & Peebles 1983; Strauss & Willick 1995). This parameter is simply related to scale-independent linear bias (Kaiser 1988), defined as the square root of the ratio of the number density variance to the mass variance $b \equiv \sigma_{\text{gal}}/\sigma_{\text{mass}} = 1/\sigma_8$. Later, σ_8 provided the standard way to normalize the theoretical power spectra of density fluctuations, determined from

first principles only up to an arbitrary multiplicative constant. A theory can be tested against the empirical data by comparing its assumed value of σ_8 with a value implied by observations. A comparison of values of σ_8 , estimated from various observations can provide an important measure of systematic errors introduced by different estimators, based on different physical effects, e.g., bulk flows, cosmic microwave background (CMB) fluctuations or gravitational lensing. Such a comparison also provides an internal consistency test for the gravitational instability theory.

To compare the $\sigma_8(z)$, estimated at high redshift, z , with its current value, estimated from observations at $z = 0$, it is necessary to take the growth of the fluctuations into account. Since all known estimates of this parameter were of the order of unity or less, until recently the linear per-

turbation theory was used for this purpose:

$$\sigma_8(0) \approx \sigma_8(z) \frac{D(0)}{D(z)} , \quad (1)$$

where D is the linear growth factor (Peebles 1980). As we show below, ignoring higher order terms in the above expression introduces a systematic error of order of 10 to 15%. With the latest improvements in the quality of observations, such effects should not be neglected and we provide a simple recipe how to take them into account. We focus on the comparison of σ_8 estimated from the peculiar velocity field at an effective redshift $z = 0$, to σ_8 implied by the CMB (effective redshift $z \approx 1100$).

Relative motions of pairs of galaxies (Feldman et al. 2003) as well as bulk flows (Pike & Hudson 2005; Sarkar et al. 2007) and velocity shear (Watkins & Feldman 2007; Feldman & Watkins 2008) measurements have been used to estimate σ_8 and Ω_m , the density of the nonrelativistic matter. In particular, from pairwise velocities we found in Feldman et al. (2003)

$$\sigma_8 = 1.13^{+0.22}_{-0.23} . \quad (2)$$

From CMB temperature fluctuations, the WMAP collaboration (Dunkley et al. 2009) found

$$\sigma_8 = 0.80 \pm 0.04 , \quad (3)$$

described as the “linear theory amplitude” (see their Table 1). The above two estimates differ only slightly, at the level of $1.5\text{-}\sigma$, and it is a success of the model that inferences by such different methods applied at two greatly different epochs are in good agreement.

In this paper we bring the two estimates even closer by taking nonlinear dynamics into account. We find that the nonlinear correction is modestly significant, given the improving accuracy in cosmology: the linear value of σ_8 , hereafter denoted σ_L , can be smaller by $\sim 10\%$ than the nonlinear value, a systematic difference that is comparable to the current statistical uncertainties. In section 2 we describe the spatial window functions used to define σ_8 . In section 3 we provide the recipe for recovering σ_L from σ . In section 4 we compare our revised σ_8 parameters to other observational estimates of σ_8 . We summarize our results in section 5.

2. Window functions

Like many cosmological experiments, measurements of cosmic flows are sensitive to a windowed integral of the matter power spectrum. In general, such an observable can be characterized as

$$Q_i = \int_0^\infty dk W(k) P_p(k) T_i^2(k) \quad (4)$$

where $P_p(k)$ gives the primordial power spectrum, k is the comoving wavenumber, $T_i^2(k)$ gives the transfer function which contains the physics of the evolution of the particular observable from the primordial spectrum. The window function, $W(k)$, describes the experimental setup (sky coverage, depth, errors etc.). This formalism describes straightforward measurements of the galaxy power spectrum, in which case $Q_i = P_{\text{gal}}(k_i)$, the CMB spectrum for which $Q_i = C_{\ell_i}$, and the amplitude of the cosmological velocity field where $Q_i = P_v(k_i)$, the velocity power spectrum. It is crucial to note that the transfer function depends implicitly upon the other cosmological parameters and hence any lack of knowledge thereof will (or at least should) translate to increased uncertainty upon the amplitude.

Bulk flow and shear measure the velocity-velocity power spectrum [or covariance, Watkins et al. (2009); Feldman et al. (2009)], whereas pairwise velocities measure the density-velocity cross-spectrum (Feldman et al. 2003). Under *linear* evolution, the density contrast is proportional to the divergence of the peculiar velocity in real space, or $\mathbf{v} \propto \mathbf{k}\rho$ in Fourier space, so these power spectra differ by powers of wavenumber k from the density power spectrum, which can be absorbed into the appropriate transfer function.

An amplitude parameter such as σ_8 is in essence a spectral observable as well. We define the *window function* $W_R(\mathbf{x}, \mathbf{x}') = W_R(\mathbf{x} - \mathbf{x}')$, normalized so that $\int d^3x W_R(x) = 1$. For our spherical top hat, $W_R(r) = 1/V$ where $V = 4\pi R^3/3$, when $r = |\mathbf{x}' - \mathbf{x}| \leq R$, 0 otherwise. Hence, the density contrast, spatially averaged over a sphere around a particular point $\mathbf{x}$ is simply

$$\delta_R(\mathbf{x}) = \int d^3x' W_R(\mathbf{x} - \mathbf{x}') \delta(\mathbf{x}') . \quad (5)$$

The ensemble average of δ_R^2 at redshift z is given

by

$$\sigma_R^2(z) \equiv \langle \delta_R^2 \rangle = \int_0^\infty \frac{dk}{k} \Delta(k, z)^2 |\tilde{W}(kR)|^2, \quad (6)$$

where

$$\tilde{W}(kR) = (3/kR) j_1(kR) \quad (7)$$

is the Fourier transform of the window function $W_R(r)$, and j_1 is a spherical Bessel function of the first kind, while

$$\Delta \equiv 4\pi P(k, z) k^3 / (2\pi)^3 \quad (8)$$

is the dimensionless power per wavenumber octave and $P(k, z)$ is the power spectrum of the mass density fluctuations. The mean square fluctuation that we would measure from the actual density contrast depends upon the actual nonlinear power spectrum, but we can analogously define the linear variance by replacing the power spectrum in the above expression with the linear spectrum, constrained to have the same amplitude for $k \rightarrow 0$ (i.e., on large scales where nonlinear evolution is negligible).

The spatial representation of the above integral is given by the expression

$$\sigma_R^2(z) = \frac{1}{V^2} \int_V d^3x d^3x' \xi(|\mathbf{x} - \mathbf{x}'|, z), \quad (9)$$

where ξ is the two-point correlation function, and $V = 4\pi R^3/3$. In Feldman et al. (2003) we have used the above equation to estimate the true present-day value of the σ_8 parameter from the PSCz survey correlation function (Hamilton & Tegmark 2002). We have also used other empirical correlation functions, derived from different surveys, as a template and we found that the resulting value of σ_8 was not sensitive to such variations; see Feldman et al. (2003).

3. Linear and nonlinear amplitudes

We use two different methods to estimate the nonlinear corrections for σ_8 , one based in perturbation theory, which allows us to express the correction as a simple analytical expression, and one using a phenomenological mapping based on conservation of pair counts and calibrated using numerical simulations, which allows us to explore the effects of changing many parameters individually.

3.1. Linear perturbation theory

Under linear evolution, the spatial and temporal dependence of clustering separates. The density perturbation $\delta(\mathbf{x}, a) = \delta\rho/\rho$ can be described as (Peebles 1980)

$$\delta(\mathbf{x}, a) = \delta^{(1)}(\mathbf{x}) D(a), \quad (10)$$

where $\delta^{(1)}$ gives the linear density perturbation field as a function of comoving spatial coordinates $\mathbf{x}$ at some fiducial time and D is the growth function, here parameterized by the scale factor a as a time coordinate (here and below we keep only the fastest-growing modes). In flat Λ CDM models the growing mode is given by (Heath 1977; Peebles 1993)

$$D(a) = \frac{5\Omega_m E(a)}{2} \int_0^a \frac{du}{u^3 E^3(u)}, \quad (11)$$

where

$$E(a) \equiv [\Omega_m a^{-3} + 1 - \Omega_m]^{1/2}, \quad (12)$$

In the early Universe, when the scale factor is small, $a \rightarrow 0$, equation (11) is well approximated by the expression

$$D(a) = a, \quad (13)$$

as in Einstein-de Sitter Universe. In the opposite limit, the cosmological constant becomes dynamically dominant and the linear growth factor will saturate at a maximum value, as gravitational clustering is balanced by the effective force of accelerated expansion. It is easy to show that in the limit $a \rightarrow \infty$, the growth factor is given by the expression

$$D_\infty = \frac{2\Gamma(\frac{2}{3})\Gamma(\frac{11}{6})}{\sqrt{\pi}} \left(\frac{\Omega_m}{1 - \Omega_m} \right)^{1/3}. \quad (14)$$

Using the above two asymptotic expressions we have found a new fitting formula for the growth factor, valid for all flat Λ CDM cosmological models:

$$D(a) = \frac{a}{[1 + (a/D_\infty)^{2.3}]^{1/2.3}}. \quad (15)$$

In Figure 1 we show that equation (15) remains within two-percent level of the exact solution (11) in both the past and the future. Note that some expressions for $D(a)$ and its logarithmic derivative, $d \log D / d \log a$, frequently quoted in the literature (Lahav et al. 1991; Carroll et al. 1992) apply only to the past and fail for $a > 1$.




Fig. 1.— The growth factor D , plotted as a function of the scale factor a for three different values of Ω_m . The exact solution (11) is represented by the solid red curve, while the dashed blue curve was derived from the approximate formula (15). All models assume zero spatial curvature, $\Omega_\Lambda = 1 - \Omega_m$. The scale factor at present equals unity. Hence the past corresponds to $a/a_0 < 1$, while the future to $a/a_0 > 1$ ($a_0 \equiv a(z=0) = 1$).

3.2. Second order perturbation theory

The perturbative solution of the equations of motion of the cosmic fluid can be written as

$$\delta(\mathbf{x}, a) = \sum_{J=1}^{\infty} \delta^{(J)}(\mathbf{x}) D^J(a) , \quad (16)$$

where $\delta^{(J)} D^J$ is the solution of order J and we assume that the linear solution is a Gaussian random field, so all its odd-order moments vanish. The mean value of the square of the above expansion is therefore given by a series of even powers of $D(a)$,

$$\langle \delta^2 \rangle = \langle [\delta^{(1)}]^2 \rangle D^2 + \langle \delta^{(2)} + 2\delta^{(1)}\delta^{(3)} \rangle D^4 + \dots \quad (17)$$

One-loop perturbative corrections to the leading order variance $\sigma_L(r)$ for power-law power spectra are given by Lokas et al. (1996) and Scoccimarro & Frieman (1996)

$$\sigma^2 = \sigma_L^2 + \beta \sigma_L^4 , \quad (18)$$

where the factor β is related to the logarithmic slope of the two-point correlation function $\gamma(r) = -d \ln \xi / d \ln r$ by the following relation:

$$\beta = 1.843 - 1.168 \gamma. \quad (19)$$

The above equations and more generally, the weakly nonlinear gravitational instability theory were confirmed by N-body simulations and by measurements of the galaxy skewness and bispectrum in redshift surveys (see e.g. Juszkiewicz et al. 1993; Scoccimarro et al. 2001; Feldman et al. 2001; Verde et al. 2002, and references therein). For non-power-law spectra, γ is a slowly varying function of scale. A convenient representation of the correlation function over scales of interest has two power laws (Hamilton & Tegmark 2002),

$$\xi(r) = q^2 (x_1^{-\gamma_1} + x_2^{-\gamma_2}) , \quad (20)$$

where $x_j = r/r_j$, $r_1 = 2.33 h^{-1} \text{ Mpc}$, $r_2 = 3.51 h^{-1} \text{ Mpc}$, $\gamma_1 = 1.72$, $\gamma_2 = 1.28$, and $q = \sigma(8 h^{-1} \text{ Mpc})/0.888$. This particular choice of set of parameters are based on the PSCz survey, and this was the choice made in the original paper, determining σ_8 from the pairwise peculiar motions of galaxies (Feldman et al. 2003). We have also considered other observational estimates of $\xi(r)$ and found that the resulting values of σ_8 and Ω_m were

unaffected. The effective slope of the correlation function is then

$$\gamma(r) \equiv -\frac{d \ln \xi(r)}{d \ln r} = \frac{\gamma_1 x_1^{-\gamma_1} + \gamma_2 x_2^{-\gamma_2}}{x_1^{-\gamma_1} + x_2^{-\gamma_2}}, \quad (21)$$

independent of the amplitude q . At $r = 8 h^{-1} \text{Mpc}$, the effective slope is $\gamma = 1.393$, for which $\beta = 0.216$. To map from σ^2 to σ_L^2 , we invert equation (18):

$$\sigma_L^2(r) = \frac{\sqrt{1 + 4\beta\sigma^2(r)} - 1}{2\beta}. \quad (22)$$

Note that for $\beta \rightarrow 0$, the above expression gives $\sigma_L^2 = \sigma^2$, as it should. In Feldman et al. (2003) we obtained $\sigma = 1.13$. Using this value with $\beta = 0.216$ in equation (22), we obtain for the central value

$$\sigma_L = 1.02, \quad (23)$$

a modest but significant decrease.

3.3. Phenomenological approach

Our other method of relating linear and nonlinear variance is non-perturbative. It uses the mapping of scale proposed in Hamilton et al. (1991), for which conservation of mass or pair counts relates a scale in the linear regime to a nonlinear ‘‘collapsed’’ scale

$$r_L^3 = \int_0^R d(r^3) [1 + \xi(r)] = R^3 [1 + \sigma^2(R)]. \quad (24)$$

The variance σ^2 is then a (nearly) universal function

$$\sigma^2(R) = g[\sigma_L^2(r_L)] . \quad (25)$$

The relation is verified and the function g identified in numerical simulations (Hamilton et al. 1991; Peacock & Dodds 1996; Smith et al. 2003).

There is an important difference between the perturbative and the phenomenological calculations. The non-perturbative mapping, given by equation (25), explicitly uses the linear expression for $D(a)$ and therefore in order to derive σ_L from σ , we have to choose specific values of the cosmological parameters; for technical details, see, for example Peacock & Dodds (1996). This is not necessary for the perturbative formula (22): to derive σ_L we only need to know σ and γ , the slope of the two-point correlation function at present.




Fig. 2.— Fully evolved variance σ plotted vs. linear variance σ_L . For comparison, the isolated red solid line shows $\sigma = \sigma_L$. Second order perturbation theory, based on Eq. (18) with $\beta = 0.216$ gives the long-dashed black line. The remaining curves are derived from the phenomenological mapping, given by Eq. (25). The differences between them show how sensitive they are to different assumptions about the cause of change in the amplitude of density perturbations. The growth induced by gravitational instability acting on density perturbations with the ‘standard’ Λ CDM initial conditions ($\Omega_m = 0.3, \Omega_\Lambda = 0.7, h = 0.7$) and the appropriate $D(a)$ factor, is shown by the blue short-dashed line. The solid red line differs in redshift or amount of evolution (and hence saturates at $\sigma_L \approx 0.95$); the dotted black line differs in Hubble constant ($h = 0.47 - 1.7$) or scale ($R = 5 - 19 \text{ Mpc}$) which changes n_{eff} and thus γ and so tracks a little differently from the others. We also show the WMAP (Dunkley et al. 2009) result ($\sigma_L = 0.80$, which maps to $\sigma = 0.85$), and that from measurements of cosmological flows ($\sigma = 1.13$ corresponding to $\sigma_L = 1.02$, with the vertical and horizontal intervals shown in long dashed black lines the range in Eqs. (26) and (27)).

In Figure 2 we show nonlinear σ plotted against the inferred linear σ (bundle of curves); for comparison the isolated solid red line shows σ_L . The perturbation theory result of equation (22) is shown by the long-dashed black line. To verify that perturbative results still make sense even at the threshold of the validity of perturbation theory, when $\sigma_L \approx 1$, we also plot the phenomenological mapping results, based on equation (25) for a variety of parameters. In this mapping we must identify why σ_L has changed, which may be from a change of fluctuation amplitude induced by the standard gravitational instability, evolution epoch, or scale; each has a slightly different effect. In general, the perturbative curve agrees well with the phenomenological results, based on the transfer function from Bardeen et al. (1986) and the Peacock & Dodds (1996) fitting. Using the Eisenstein & Hu (1998) transfer function and the Smith et al. (2003) fitting make very little difference in the results except when the Hubble constant or scale changes. This occurs because a change in scale substantially changes the value of γ and so of β . This is yet another confirmation of the reliability of our perturbative calculations. It also shows that the PSCz $\xi(r)$, assumed in the perturbative calculation, agrees well with the ‘standard’ Λ CDM $\xi(r)$, assumed in the phenomenological mapping.

In the phenomenological Λ CDM relation, the nonlinear signal range

$$\sigma = 1.13^{+0.22}_{-0.23}, \quad (26)$$

maps to

$$\sigma_L = 1.02^{+0.16}_{-0.18}. \quad (27)$$

Note that since σ is steeper than σ_L , the range in σ_L is somewhat narrower than the one in σ . These values and the resulting limits are also shown in Fig. 2.

More generally, the existing descriptions and ansatzes of nonlinear clustering were created to describe the evolution of clustering from the past through to the present-day; they are not necessarily adequate representations of future clustering, even in the very near future. In Figure 3 we show the value of $\Delta^2(k, z)$, the mean square of the dimensionless density fluctuation at various epochs using both the Peacock & Dodds (1996) mapping and the Smith et al. (2003) formula; the former

becomes time independent whereas the actual evolution will concentrate more and more matter in smaller haloes over time. We also see this in Figure 4 which shows the value of both σ_8 and σ_1 as a function of expansion factor. The latter scale of $1h^{-1}$ Mpc is nonlinear today and therefore already shows a difference between linear evolution and the different nonlinear formulae, whereas the canonical $8h^{-1}$ Mpc scale is just going nonlinear today.




Fig. 3.— The linear (short-dashed black lines) and nonlinear values of the dimensionless $\Delta^2(k)$ as obtained from the nonlinear ansatz of Peacock & Dodds (1996, long-dashed red lines) and from Smith et al. (2003, solid blue lines), for expansion factors $a = 1/4, 1/2, 1, 2, 4$, and 8 (where the present is $a = 1$).

These differences arise partly from the fact the various fitting formulae did not attempt to reproduce future clustering, but moreover, from the well-known fact that we live at a special time: in Λ CDM, linear clustering will “saturate” soon. Hence, any prediction such as that of Peacock & Dodds (1996) based on a mapping of the linear spectrum will also necessarily saturate. Moreover, the various densities are currently evolving very rapidly. For example, at $a = 1/2$, Ω_m was approximately what Ω_Λ is today; at $a = 2$, Ω_m will be approximately what Ω_b is today. This also gives us some insight into




Fig. 4.— The value of σ_8 (lower set of curves) and σ_1 (upper set of curves) as a function of expansion factor, for linear theory (short-dashed black lines), Peacock & Dodds (1996, long-dashed red lines) and from Smith et al. (2003, solid blue lines).

the underlying (rather than just practical) limitations of these methods. The mapping proposed by Peacock & Dodds (1996) essentially applies equally to all scales. However, in an evolved Λ CDM universe, large-scale dynamics are dominated by the accelerated expansion and small scales by highly nonlinear stable clustering, which is captured somewhat better in the halo model.

4. Discussion

We will now compare different estimates of σ_8 .

CMB. The σ_8 is estimated from the CMB fluctuations, observed at $z \approx 1100$ and then linearly extrapolated to the present era. The CMB itself measures a complicated (albeit linear) functional of the power spectrum, and because of degeneracies in the determination of the CMB power spectrum from the cosmological parameters, the value of σ_8 from the CMB depends on details of the data-analysis procedure, in particular on the assumed Bayesian priors on the cosmological parameters considered. The “recommended” value for a flat Λ CDM

model (<http://lambda.gsfc.nasa.gov>) is $\sigma_8 = 0.80 \pm 0.04$. However, considering different models and priors on those parameters can give variations over $0.7 \lesssim \sigma_8 \lesssim 0.9$.

Cosmic flows. Observations of peculiar velocities of galaxies provide an estimate of σ_8 at the present, nonlinear regime. Masters et al. (2006) estimated $\sigma_8 \Omega_m^{0.6} = 0.52 \pm 0.06$ using the peculiar velocities from a sample of clusters in the SFI++ survey; Zaroubi et al. (2002) using density-density and velocity-velocity comparisons found $\sigma_8 \Omega_m^{0.6} = 0.45 \pm 0.05$ both of which translate to $\sigma_8 \gtrsim 1$. At scales of $100 h^{-1}$ Mpc, Lavaux et al. (2008) found $\sigma_8 = 1.72 \pm 0.56$ from 2MASS, although they only directly measure the velocity field within $30 h^{-1}$ Mpc, on which scales they estimate a somewhat lower fluctuation amplitude. As mentioned above, pairwise velocity analysis (Feldman et al. 2003) estimates a present day nonlinear value of $\sigma_8 = 1.13^{+0.22}_{-0.23}$. Recent bulk flow measurements using the best available peculiar velocity surveys (Watkins et al. 2009) require the nonlinear $\sigma_8 > 1.11$ (0.88) at a 95% (99%) confidence level. Thus, flow measurements give higher values of σ_8 than the CMB, as expected from the ideas discussed above. We expect more precise measurements of σ_8 from cosmic flows in the near future. The new surveys are deeper, denser, have better sky coverage (Masters et al. 2006; Springob et al. 2007, 2009), and we have improved our understanding of the distance indicators needed to extract the peculiar velocities (Feldman et al. 2003; Radburn-Smith et al. 2004; Pike & Hudson 2005; Sarkar et al. 2007; Watkins & Feldman 2007; Feldman & Watkins 2008).

Other cosmological probes. For comparison, we provide a table, where apart from the above estimates of σ_8 , we also show the values, derived from Ly α observations (Tytler et al. 2004); cluster number density measurements (Vikhlinin et al. 2009); weak lensing measurements of cosmic shear (Benjamin et al. 2007); the Sunyaev-Zeldovich effect using the ACBAR receiver (Reichardt et al. 2009), and from the galaxy clustering power spectrum (Cole et al. 2005; Tegmark et al. 2004; Eisenstein et al. 2005). All of these measurements fall in the range $0.7 < \sigma_8 < 1.2$ with the most likely values in the interval $0.8 < \sigma_8 < 0.95$.

Table 1: σ_8 from various estimators

Method	Parameter	value
CMB ¹	σ_L	0.80 ± 0.04
LY- α ²	σ_L	0.85 ± 0.02
Cosmic Shear ³	σ_L	0.84 ± 0.05
Clusters ⁴	σ_L	0.75 ± 0.01
SZ (ACBAR) ⁵	σ_8	$0.94^{+0.03}_{-0.04}$
Galaxies ⁶	σ_8^{gal}	0.92 ± 0.06
Flows		
Pairwise velocities ⁷	σ_8	$1.13^{+0.22}_{-0.23}$
Bulk flow ⁸	σ_8	> 1.11 (0.88) at 95% (99%) CL

¹Dunkley et al. (2009); ²Tytler et al. (2004); ³Benjamin et al. (2007); ⁴Vikhlinin et al. (2009); ⁵Reichardt et al. (2009); ⁶Cole et al. (2005); Tegmark et al. (2004); Eisenstein et al. (2005); ⁷Feldman et al. (2003); ⁸Watkins et al. (2009)

5. Conclusions

We have presented a formalism to calculate σ_8 , the amplitude of cosmological density fluctuations on scales of $8 h^{-1}$ Mpc, that incorporates the non-linear evolution of the parameter. Estimates of σ_8 depend directly on the epoch and scale of the surveys used. When using deep, high-redshift surveys (CMB) that estimate σ_8 in the linear regime, the results suggest systematically lower values of σ_8 . When analyzing shallow, local data (peculiar velocities) which are affected by nonlinearities, we get higher σ_8 . The results from other cosmological probes, shown in Table 1, show a similar trend. When results from deep surveys are being corrected for this effect, most estimates from various independent surveys on all scales agree better with each other: the systematic differences are reduced.

Quantitatively, our main result here is the reduction in the value of σ_8 , derived from the observed mean pairwise velocity of galaxies. The original estimate, as we have discussed earlier, is

$$\sigma_8 = 1.13^{+0.22}_{-0.23}. \quad (28)$$

After the correction based on the second order perturbation theory, this becomes

$$\sigma_L = 1.02^{+0.16}_{-0.18}. \quad (29)$$

Somewhat more cumbersome non-perturbative methods give identical results, bringing late-time

estimates of σ_8 close to high-redshift measurements, which appear in Table 1.

Acknowledgements: We would like to thank the anonymous referee for her/his constructive critical remarks. RJ was supported by the Polish Ministry of Science grant NN203 2942 34 and an INTAS grant No. 06-1000017-9258. HAF has been supported in part by a grant from the Research Corporation, by an NSF grant AST-0807326 and by the National Science Foundation through TeraGrid resources provided by the NCSA. AHJ was supported by STFC in the UK.

REFERENCES

- Bardeen, J. M., Bond, J. R., Kaiser, N., & Szalay, A. S. 1986, ApJ, 304, 15
- Benjamin, J., et al. 2007, MNRAS, 381, 702
- Carroll, S. M., Press, W. H., & Turner, E. L. 1992, ARAA, 30, 499
- Cole, S., et al. 2005, MNRAS, 362, 505
- Davis, M., & Peebles, P. J. E. 1983, ApJ, 267, 465
- Dunkley, J., et al. 2009, ApJS, 701, 1804
- Eisenstein, D. J., & Hu, W. 1998, ApJ, 496, 605
- Eisenstein, D. J., et al. 2005, ApJ, 633, 560
- Feldman, H., et al. 2003, ApJ, 596, L131

- Feldman, H. A., Frieman, J. A., Fry, J. N., & Scoccimarro, R. 2001, PRL, 86, 1434
- Feldman, H. A., & Watkins, R. 2008, MNRAS, 387, 825
- Feldman, H. A., Watkins, R., & Hudson, M. J. 2009, ArXiv e-prints
- Hamilton, A. J. S., Kumar, P., Lu, E., & Matthews, A. 1991, ApJ, 374, L1
- Hamilton, A. J. S., & Tegmark, M. 2002, MNRAS, 330, 506
- Hamilton, A. J. S., & Tegmark, M. 2002, MNRAS, 330, 506
- Heath, D. J. 1977, MNRAS, 179, 351
- Juszkiewicz, R., Bouchet, F. R., & Colombi, S. 1993, ApJ, 412, L9
- Kaiser, N. 1988, MNRAS, 231, 149
- Lahav, O., Lilje, P. B., Primack, J. R., & Rees, M. J. 1991, MNRAS, 251, 128
- Lavaux, G., Tully, R. B., Mohayaee, R., & Colombi, S. 2008, ArXiv e-prints
- Lokas, E., Juszkiewicz, R., Bouchet, F. R., & Hivon, E. 1996, Astrophys. J., 467, 1
- Masters, K. L., Springob, C. M., Haynes, M. P., & Giovanelli, R. 2006, ApJ, 653, 861
- Peacock, J. A., & Dodds, S. J. 1996, MNRAS, 280, L19
- Peebles, P. J. E. 1980, The large-scale structure of the universe (Princeton, N.J., Princeton University Press, 1980)
- . 1993, Principles of physical cosmology (Princeton, N.J., Princeton University Press, 1980)
- Pike, R. W., & Hudson, M. J. 2005, ApJ, 635, 11
- Radburn-Smith, D. J., Lucey, J. R., & Hudson, M. J. 2004, MNRAS, 355, 1378
- Reichardt, C. L., et al. 2009, ApJ, 694, 1200
- Sarkar, D., Feldman, H. A., & Watkins, R. 2007, MNRAS, 375, 69
- Scoccimarro, R., Feldman, H. A., Fry, J., & Frieman, J. 2001, ApJ, 546, 652
- Scoccimarro, R., & Frieman, J. A. 1996, ApJS, 105, 37
- Smith, R. E., et al. 2003, MNRAS, 341, 1311
- Springob, C. M., Masters, K. L., Haynes, M. P., Giovanelli, R., & Marinoni, C. 2007, ApJS, 172, 599
- . 2009, ApJS, 182, 474
- Strauss, M. A., & Willick, J. A. 1995, Phys. Rep., 261, 271
- Tegmark, M., et al. 2004, ApJ, 606, 702
- Tytler, D., et al. 2004, ApJ, 617, 1
- Verde, L., et al. 2002, MNRAS, 335, 432
- Vikhlinin, A., et al. 2009, ApJ, 692, 1033
- Watkins, R., & Feldman, H. A. 2007, MNRAS, 379, 343
- Watkins, R., Feldman, H. A., & Hudson, M. J. 2009, MNRAS, 392, 743
- Zaroubi, S., Branchini, E., Hoffman, Y., & da Costa, L. N. 2002, MNRAS, 336, 1234